

Hormonal regulation of hippocampal dendritic morphology and synaptic plasticity

Peter R. Moulton & Jenni Harvey

To cite this article: Peter R. Moulton & Jenni Harvey (2008) Hormonal regulation of hippocampal dendritic morphology and synaptic plasticity, Cell Adhesion & Migration, 2:4, 269-275, DOI: [10.4161/cam.2.4.6354](https://doi.org/10.4161/cam.2.4.6354)

To link to this article: <https://doi.org/10.4161/cam.2.4.6354>



Copyright © 2008 Landes Bioscience



Published online: 01 Oct 2008.



Submit your article to this journal [↗](#)



Article views: 130



Citing articles: 39 View citing articles [↗](#)

Special Focus: Recent Advances in the Cellular and Molecular Mechanisms Underlying Synaptic Plasticity

Hormonal regulation of hippocampal dendritic morphology and synaptic plasticity

Peter R. Moulton and Jenni Harvey*

Neurosciences Institute; Division of Pathology and Neuroscience; Ninewells Hospital and Medical School; University of Dundee; Dundee, UK

Key words: synaptic plasticity, leptin, estrogen, insulin, hippocampus, LTD, LTP

The peripheral functions of hormones such as leptin, insulin and estrogens are well documented. An important and rapidly expanding field is demonstrating that as well as their peripheral actions, these hormones play an important role in modulating synaptic function and structure within the CNS. The hippocampus is a major mediator of spatial learning and memory and is also an area highly susceptible to epileptic seizure. As such, the hippocampus has been extensively studied with particular regard to synaptic plasticity, a process thought to be necessary for learning and memory. Modulators of hippocampal function are therefore of particular interest, not only as potential modulators of learning and memory processes, but also with regard to CNS driven diseases such as epilepsy. Hormones traditionally thought of as only having peripheral roles are now increasingly being shown to have an important role in modulating synaptic plasticity and dendritic morphology. Here we review recent findings demonstrating that a number of hormones are capable of modulating both these phenomena.

Introduction

It is well documented that hormones play a vital role in the regulation and processing of numerous biochemical pathways throughout the body. Until recently, it was thought that hormonal regulation of such processes was restricted largely to the periphery. However, in recent years it has emerged that a range of hormones can be found within the CNS along with complimentary receptors capable of translating hormonal signals into biological actions. These hormone receptor complexes have been shown to have a wide variety of modulatory actions at the neuronal level, no more so than in the hippocampus. Since the early 1950's the hippocampus has been recognised to play a fundamental role in certain forms of learning and memory. It is an area where synaptic plasticity, thought to be the cellular correlate of learning and memory, has been extensively investigated and therefore hormonal modulation of neuronal func-

tion in the hippocampus is of great interest and has wide spread implications. Although a number of hormonal systems have been shown to modulate hippocampal function, the largest bodies of work center around estrogens, leptin and insulin. These are exciting and rapidly expanding fields with regard to hippocampal modulation and for this reason we have focused this review on some of the recent work investigating the modulation of hippocampal dendritic morphology and synaptic plasticity mediated by these hormones.

Leptin

Leptin is a 167 amino acid protein produced almost exclusively in adipose tissue and is found to circulate in the plasma at levels relative to body fat tissue.^{1,2} It is well documented that leptin acts on a number of peripheral tissues, but there is strong evidence that leptin can cross the blood brain barrier and act upon targets within the CNS. Moreover, leptin mRNA has been described in a number of brain regions giving rise to the possibility leptin may be synthesised and released from within the CNS itself.³ Interest in leptin initially focused on its centrally mediated weight-reducing effects and on the potential for using the leptin/leptin receptor axis to develop therapeutic drugs to treat obesity.

The most widely documented functions of leptin within the CNS are found to be in the hypothalamus, where leptin is involved in regulating energy homeostasis,⁴ bone formation,⁵ reproduction⁶ and the hypothalamic-pituitary-adrenal axis.⁷ It has emerged recently however that the functions of leptin within the CNS are not restricted solely to the hypothalamus, but that leptin is involved in a diverse array of processes across the CNS, most notably in the hippocampus.⁸⁻¹⁰

The leptin receptor (Ob-R) was first cloned in 1995 from the mouse choroid plexus.¹¹ Since then, six leptin receptor isoforms, generated by alternate splicing of the *db* gene have been identified in rodents.¹² The six leptin receptor isoforms are termed Ob-Ra to Ob-Rf and have identical extracellular, N-terminal domains. The differences between the isoforms occur at the intracellular C-terminal domains. All isoforms except Ob-Re have a 34 amino acid transmembrane domain. Ob-Re is thought to act as a soluble receptor within the plasma facilitating leptin transport. The membrane spanning isoforms of the leptin receptor are divided into two categories based upon the length of the c-terminal domain. Ob-Ra, c, d and f

*Correspondence to: Jenni Harvey; University of Dundee; Neurosciences Institute; Ninewells Hospital; Dundee, Tayside, DD1 9SY UK; Email: j.z.harvey@dundee.ac.uk

Submitted: 02/14/08; Accepted: 05/28/08

Previously published online as a *Cell Adhesion & Migration* E-publication: <http://www.landesbioscience.com/journals/celladhesion/article/6354>

have a shorter intracellular domain and are thus termed short forms whilst Ob-Rb has a larger intracellular domain and is thus termed the long form. Although membrane bound short forms of the receptor are able to activate certain signalling cascades, their major role is likely to be related to leptin internalization and degradation.¹³ Another possibility is that they are involved in leptin clearance or receptor mediated transport into the brain¹⁴ (reviewed in ref. 15). The longer C terminus of Ob-Rb contains 302 residues in which reside a number of motifs which facilitate the initiation of various downstream signalling cascades.

In line with leptin's reported regulation of hypothalamic function, levels of leptin Ob-Rb mRNA are found to be particularly high in specific regions of the hypothalamus, that are involved in regulating food intake and body weight.¹⁶⁻¹⁹ To date, a number of studies have found high levels of Ob-Rb mRNA expression in several CNS regions including the amygdala, cerebellum, brainstem, substantia nigra and also the hippocampus.¹⁸⁻²¹ As well as in the hypothalamus, particularly high levels (relative to overall CNS expression) of Ob-R mRNA and immunoreactivity are apparent in hippocampal CA1 and CA3 regions as well as in the dentate gyrus.^{19,20,22} In primary hippocampal cultures, leptin receptors have been shown to be expressed on pyramidal neurons and on glial cells.²³ Moreover, leptin receptor immunostaining is found at axonal and somato-dendritic regions and colocalises with the synaptic marker, synapsin I,²³ suggesting it is localised to synapses. Again, the presence and indeed the high level of expression of both Ob-Rb mRNA and Ob-Rb itself, suggests a role for leptin in the regulation of hippocampal function. It is also evident that Ob-Rb is absent in the choroid plexus, meninges, and also surrounding blood vessels, yet a probe recognizing all known forms of the leptin receptor indicates expression in these areas.¹⁸ This suggests expression of the short isoforms, adding support to the idea that they may be involved in transport of leptin into the brain. As more evidence emerges it is becoming apparent that leptin plays a major role in the regulation of hippocampal synaptic transmission, as well as synaptic plasticity and dendritic morphology. Although some of the studies above have focused on the long form Ob-Rb receptor, it should be remembered that many have looked at general Ob-R mRNA or Ob-R expression. It is likely that effects in these brain regions are mediated by Ob-Rb, however in the absence of isoform specific antagonists we cannot rule out the involvement of other isoforms.

The ability to regulate synaptic strength is a critical feature of the CNS and is thought to underlie processes such as learning and memory. Long-term potentiation (LTP) and long-term depression (LTD) have been studied extensively in the hippocampus to understand the mechanisms of synaptic plasticity.^{24,25} Recent evidence has shown that leptin not only regulates hippocampal synaptic transmission per se, but can also regulate different forms of hippocampal synaptic plasticity. Strong evidence that leptin is able to regulate both hippocampal LTP and LTD comes from studies of rodents with Ob-R mutations (either *db/db* mice or *fafa* rats) exhibiting both reductions in LTP and LTD²⁶ as recorded in the CA1 region of acute hippocampal slices. Moreover these rodent strains also show impaired spatial memory in a variety of behavioural tests.^{26,27} Conversely, direct administration of leptin into the hippocampus has been shown to improve learning and memory^{28,29} as well as to facilitate LTP.³⁰

NMDA receptors contribute little to basal synaptic transmission, but are crucial for the induction of certain forms of synaptic plasticity. Leptin has been shown to selectively enhance NMDA receptor-mediated synaptic currents at CA1 synapses in acute hippocampal slices.³¹ Furthermore, exposure of acute hippocampal slices to leptin promotes the conversion of short-term potentiation (STP; induced by primed burst stimulation) into LTP.³¹ As part of this same study, activation of leptin receptors increased NMDA receptor mediated currents in *Xenopus* oocytes expressing recombinant NMDA receptor channels. It seems therefore that leptin is able to modulate hippocampal LTP via facilitation of NMDA receptor function. Changes in the morphology and/or the density of dendritic spines have been shown to contribute to enhanced synaptic efficacy following hippocampal LTP.³² In this respect, it has recently been shown that, in an NMDA receptor dependent manner, leptin rapidly enhances the motility and density of dendritic filopodia in cultured hippocampal neurones and enhances the number of synapsin I positive puncta seen within these cells.³³

The signalling cascades activated downstream of leptin receptors are much the same as other members of the class I cytokine receptor superfamily and utilise associated janus tyrosine kinases (JAKs). Ligand binding to the leptin receptor results in transphosphorylation of JAK2 and phosphorylation of receptor tyrosine residues, Tyr¹¹³⁸ and Tyr⁹⁸⁵, on the intracellular domain. This leads to the activation of various downstream signalling molecules, including the STAT (signal transducers and activators of transcription) family of transcription factors, insulin receptor substrate (IRS) proteins, phosphoinositide 3-kinase (PI3-kinase) and adaptor proteins associated with Ras-Raf-MAPK (mitogen-activated protein kinase) signalling cascade.

Several studies have examined the cell signalling cascades coupling leptin receptor activation to the modulatory effects on hippocampal synaptic plasticity. The facilitation of NMDA receptor mediated currents was found to be dependent on both PI3 kinase and the MAPK, ERK.³¹ Conversely, the rapid increase in motility and density of dendritic filopodia and subsequent enhancement of the density of synaptic labelling was shown to be independent of PI3 kinase, but was dependent on MAPK.³³ Several lines of evidence suggest different localization of different NR2, NMDA receptor subunits within the hippocampus; NR2A containing NMDA receptors being predominantly synaptic whilst NR2B are predominantly extrasynaptic.^{34,35} The rapid increase in motility and density of dendritic filopodia induced by leptin is dependent upon synaptic, i.e., NR2A containing, NMDA receptors as opposed to NR2B. Interestingly, a number of studies suggest that activation of NR2A containing NMDA receptors is a pre-requisite for the induction of hippocampal an cortical LTP, whereas NR2B containing NMDA receptors are activated during LTD.^{36,37}

As well as modulating hippocampal LTP, leptin has been recently shown to induce a novel form of hippocampal LTD, induced under conditions of increased excitability.³⁸ Leptin induced LTD shares many characteristics with synaptically induced NMDA receptor-dependent LTD, i.e., NMDA receptor dependence, postsynaptic expression and mGluR-independence. The signalling mechanisms underlying this form of LTD do differ from that of low frequency stimulation (LFS)-induced LTD though; leptin-induced LTD is negatively regulated by PI3-kinase and serine/threonine phosphatases 1/2A.³⁸

Insulin

Insulin is a 6 KDa, protein which is synthesized in significant quantities by pancreatic β -cells. Appropriate stimulation of β -cells causes secretion of insulin by exocytosis which then diffuses into islet capillary blood. Binding of insulin to the insulin receptor then regulates the uptake of glucose from the circulation. It is well established that insulin regulates blood glucose levels in this way and, much like leptin, for many years its actions were believed to be restricted to the periphery since insulin was believed to be incapable of crossing the blood brain barrier. It was later discovered that insulin receptors were present in the CNS, originally identified through *in vitro* binding studies.³⁹⁻⁴¹ Further studies identified insulin receptors in both neurons and glia throughout the rat brain⁴² and more specifically, the β subunit colocalised with synaptophysin in cultured hippocampal neurons as well as being enriched in synaptosome and PSD fractions.⁴³ Moreover, insulin was later shown to in fact be capable of crossing the blood brain barrier⁴⁴ and there is evidence to suggest it may even be synthesized and released by neurons.⁴⁵⁻⁴⁷ This mounting evidence all implicates a functional role for insulin within the CNS. Indeed, it is now established that insulin influences several CNS functions, including hypothalamic driven feeding,⁴⁸ excitatory and inhibitory synaptic transmission⁴⁹⁻⁵¹ and neuronal survival.⁵² Moreover, dysfunctional insulin signalling is thought to contribute to the development of neurodegenerative disorders such as Alzheimer's disease.⁵³

Insulin binding to the insulin receptor leads to intracellular recruitment of insulin receptor substrates and activation of PI3K signalling.⁵⁴⁻⁵⁶ In addition, recruitment of the adaptor protein SHC to the insulin receptor leads to activation of the ERK 1/2 pathway.^{54,57} It is thought that insulin activation of ERK 1/2 may underlie changes in synaptic plasticity and hippocampal dependent learning and memory.⁵⁸⁻⁶⁰ A role for PI3K is also emerging, as PI3K signalling has been implicated in the induction and maintenance of LTP^{61,62} and specific forms of LTD.⁶³

There is a growing body of evidence to suggest that insulin, like leptin, can influence synaptic plasticity in the CNS, and more specifically in the hippocampus. Brief application of insulin to hippocampal slices evokes LTD of excitatory synaptic transmission.⁶⁴⁻⁶⁶ Similar to both synaptically induced LTD (evoked by LFS) and leptin-induced LTD, LTD induced by insulin is dependent upon NMDA receptor activation, is expressed postsynaptically and is mGluR-independent. A number of downstream signalling pathways have been implicated in insulin-induced LTD, including PI3K and PKC.^{65,66} Recent studies suggest that the mechanism involves tyrosine phosphorylation of the GluR2 AMPA receptor subunit⁶⁷ and internalization of GluR2 containing AMPA receptors.^{64,65}

Contrastingly, cell surface expression of the GluR1 AMPA receptor subunit increases following exposure of hippocampal neurons to insulin.^{64,68} Interestingly, increases in cell surface GluR1 expression are also thought to underlie hippocampal LTP.⁶⁹ Therefore, like leptin, insulin has the unusual capacity to increase as well as decrease synaptic efficacy. Insulin also enhances NMDA receptor mediated currents in hippocampal neurons⁷⁰ and promotes an increase in the cell surface density of NMDA receptors.⁵¹ This potentiation of NMDA currents is dependent upon both tyrosine phosphorylation and PKC.^{70,71} As these effects are likely to lower

the threshold for the induction of plasticity it is not surprising that insulin shifts the frequency response curve of synaptic plasticity to the left.⁶⁶ Stimulation of rat hippocampal slices with insulin also rapidly increases the protein expression of PSD-95 in the CA1 area in a PI3K dependent manner.⁷² As PSD-95 functions as a scaffold protein, assembling a specific set of signalling proteins around the NMDA receptor,⁷³ regulation of PSD-95 might provide an additional molecular mechanism by which insulin could modulate hippocampal synaptic plasticity.

To date, there is little evidence to suggest that insulin, via activation of insulin receptors, modulates hippocampal dendritic morphology. However insulin like growth factor 1 (IGF-1) does promote increased dendritic branching in cortical, pyramidal neurons⁷⁴ and furthermore cortical neurons from IGF1^{-/-} mice exhibit a reduction in dendritic spine density.⁷⁵ Given that insulin has a high affinity for and can signal via IGF-1 receptors, it raises the possibility that insulin can also modulate dendritic morphology. Furthermore, there is evidence to suggest that insulin modulates dendritic growth and neurite outgrowth in a number of brain regions and cell types (reviewed in refs. 52 and 76). Modulation of hippocampal dendritic morphology by insulin is clearly an important area of research which needs to be pursued.

Estrogens

Early studies of estrogens focussed on reproduction and estrogenic actions upon reproductive target organs. As with so many hormones, it has become evident that estrogens acts to influence many other behaviors including modulating forms of learning and memory as well as other cognitive functions (reviewed in ref. 77).

Estrogenic effects within the CNS and more specifically the hippocampus are mediated by estrogens receptors (ERs). There is a wide body of evidence to support estrogen signalling mediated by both nuclear and membrane/synaptically located ERs.^{78,79} Genomic signalling (via nuclear receptors) involves nuclear estrogen-receptor complexes binding estrogen response elements (EREs) in the DNA and regulating gene transcription. Non-genomic signalling (via membrane bound receptors) involves membrane bound receptors activating intracellular cascades upon estrogens binding. To date, evidence suggests that both these methods of signalling occur in a number of hippocampal cell types shown to mediate estrogenic effects.

Estrogen receptor α (ER α) and estrogen receptor β (ER β) are the best characterized estrogen receptors and both are expressed throughout the hippocampal formation.⁸⁰⁻⁸² Both receptor subtypes have the potential to mediate nuclear as well as membrane associated effects.⁸³ Other receptor systems associated with estrogens exist in the CNS, but to date remain poorly understood.⁸⁴

The primary source of CNS estrogens are derived from the periphery. In females, estrogen synthesis occurs in the ovaries where testosterone is converted into estradiol within the granulosa cells. This same process occurs in the gonads of males, but to a much lesser extent. Evidence suggests that, like insulin and leptin, estrogens can be transported across the blood brain barrier allowing it to mediate processes within the CNS. Local synthesis of estrogens within the CNS is also a possibility, specifically within the hippocampus. It has been demonstrated *in vitro* that hippocampal neurons can synthesise estradiol,⁸⁵⁻⁸⁸ raising the possibility that locally produced estradiol may mediate estrogenic sensitive hippocampal effects.

However, evidence for local production of estrogens to date stem, only from studies of cultured cells or isolated tissue and as of yet no clear evidence exists that estrogens are synthesised *in vivo* in the hippocampus.

The distribution of ERs within the hippocampus is contentious. ER α and ER β have been reported throughout all principle cells of the hippocampus as well as in glial cells and different types of GABAergic interneuron (reviewed in ref. 89). Early studies suggested that ERs were principally expressed in GABAergic cells and that estrogens regulate hippocampal activity via modulation of inhibition. Electron microscopy studies have also identified ER α and ER β in numerous sub-cellular locations,^{90,91} but limitations of current antibodies and the fact that ERs have been shown to be differentially regulated at different stages of the oestrous cycle may go some way to explaining the above discrepancies. What is without doubt is that ERs are expressed throughout the hippocampus, although as to what extent remains contentious and that they mediate the ability of estrogens to modulate hippocampal dendritic morphology and synaptic plasticity.

Despite conflicting data as to the exact location and expression levels of ERs in the hippocampus, it is clear that estrogens have a profound effect upon dendritic morphology, particularly at the synaptic level. Ovariectomy in rats decreases the density of dendritic spines in the CA1 region of the hippocampus; an effect that can be reversed by estradiol replacement.^{92,93} Furthermore, estradiol increases spine density in rat primary hippocampal neurons in culture.⁹⁴ A similar change in spine and synapse density also occurs within the natural oestrous cycle. Rats in late proestrus have a 30% higher density of spines in the CA1 than in late oestrus.⁹⁵ Proestrus rats also have a higher proportion of "mushroom" as opposed to "stubby" spines, with evidence suggesting "mushroom" shaped spines to be a more mature and stronger subset of spines.^{95,96} Thus as well as increasing spine size, estradiol may play a role in the maturation of dendritic spines.

Consistent with estradiol increasing spine density, estradiol also increases excitatory synaptic contact when observed at the ultrastructural level.⁹⁷ Increases in spine density correlate with an increase in the number of pre-synaptic boutons; estradiol increases the number of synapses by 25%.⁹³ This data, obtained through electron microscopy, provides direct evidence of an increase in the number of synapses in response to estradiol. More recent evidence also shows increases in the molecular components of both postsynaptic spines and presynaptic boutons in response to estradiol. In the CA1 region, levels of synaptophysin, syntaxin, spinophilin and PSD-95 all increase in response to estradiol exposure in both ovariectomised rats^{98,99} and in cultured hippocampal neurons.^{94,98,100,101}

As far back as 1980 it was reported that estradiol could acutely modulate hippocampal excitability. Electrophysiological studies have shown acute modulation of synaptic events, presumably via membranous ERs. Application of estradiol to hippocampal slices increased the amplitude of synaptically evoked population spikes in the CA1 region of the hippocampus.¹⁰² Subsequently 17 β estradiol was reported to increase the CA1, dendritic fEPSP,¹⁰³⁻¹⁰⁵ a finding more recently extended to both the dentate gyrus and CA3 regions of the hippocampus.¹⁰⁶ The studies outlined above highlight gross effects on a population of neurons, but single cell recordings largely mirror these results.^{107,108} Enhancement of AMPA receptor mediated synaptic transmission by estradiol is independent of NMDA

receptor activation, however, NMDA receptor mediated EPSPs can also be facilitated.¹⁰³

Given the marked effects of estradiol on basal synaptic transmission and dendritic morphology in the hippocampus it is not surprising that estradiol also acts as a modulator of hippocampal synaptic plasticity. Indeed, the magnitude of LTP elicited by brief trains of high frequency stimulation (HFS) is greater in female rats during proestrus.¹⁰⁹ Furthermore, ovariectomised rats treated with estradiol were found to show facilitation of hippocampal LTP,¹¹⁰ an enhancement of hippocampal LTD¹¹¹ as well as a reduction in the frequency threshold for the induction of hippocampal LTD.¹¹² These effects of estradiol may also be dependent upon age as in slices from 3–4 week old rats, estradiol has been reported to suppresses hippocampal LTP.¹¹³

As previously discussed, the ability of synapses to undergo long-lasting alterations in efficacy, via the process of synaptic plasticity, is thought to be necessary for learning and memory.²⁴ The hippocampus is a major mediator of spatial learning and memory and it has recently been demonstrated that learning induces LTP in the hippocampus *in vivo*.¹¹⁴ In keeping with this, as well as facilitating LTP and LTD, estradiol enhances the performance of rodents in spatial learning tasks.¹¹⁵⁻¹¹⁸ Furthermore, in a manner similar to studies on receptor distribution, spine modulation and synaptic physiology, the performance of rodents in memory tasks fluctuates across the oestrus cycle and shifts in the preferred learning strategy have been observed.¹¹⁵

It is unclear whether the actions of estrogens discussed above are mediated via nuclear or membranous ERs. Nuclear estradiol binding and ER α immunoreactivity has been seen in hippocampal GABAergic interneurons¹¹⁹ with nuclear ER β seen in astrocytes¹²⁰ in the hippocampus. This raises the possibility that some estrogenic effects in the hippocampus, especially those involving mediation of GABAergic cells may occur via a genomic effect. Nuclear estradiol binding has also been seen in the basal forebrain which projects to the hippocampal formation.¹²¹ However, it should be noted that there is no direct evidence providing a function for these nuclear hippocampal receptors in the forebrain.

As alluded to earlier, non genomic signalling through ERs has been identified outside the nucleus. The extranuclear receptors which mediate this signalling act by coupling to G proteins, growth factor receptors, and intracellular kinases signalling both rapid hormonal actions and also more delayed effects on gene transcription by activating intracellular signal cascades.^{79,122} These receptors are well positioned to process many of the effects described above e.g., on axonal terminals and also on dendritic spines.^{90,123} Many of the estrogenic effects described above occur too quickly to be attributable to gene transcription and so it is more likely that they are mediated via non nuclear receptors. Furthermore some of the effects described are post-transcriptional, suggesting effects on protein translation.^{98,101}

It is clear that the available evidence is inconclusive as to which receptor systems mediate many of the modulatory effects of estrogens on hippocampal dendritic morphology and synaptic plasticity. However, a recent study has shown that selective agonists for ER β increase key synaptic proteins, an effect absent in ER β knockout mice.¹²⁴ Furthermore, this same study demonstrated increased dendritic branching, increased spine density and improved




Figure 1. Simplified, schematic diagram showing how leptin, estrogen and insulin signalling may converge upon the same signalling pathways in order to modulate hippocampal synaptic plasticity and dendritic morphology. Upon appropriate ligand binding either receptor is capable of activating either the Ras/Raf/MEK or the PI3K/Akt signalling cascades, both of which have been implicated in various forms of hippocampal synaptic plasticity.

performance in hippocampal dependent memory tasks in response to a specific ER β agonist and not to an ER α agonist.

It is likely that hippocampal function is altered via a combination of genomic and non-genomic effects depending upon the specific subregion or cell type. It is also possible that some effects may require a genomic and non-genomic component and that these two signalling mechanisms act together to mediate a common goal.

Concluding Remarks

Hippocampal function and modulation of hippocampal function is an area of great interest and an area that has been extensively investigated. Evidence is mounting that hormonal regulation of hippocampal cellular events such as synaptic plasticity plays a vital role in the way the hippocampus functions. As the list of hormonal modulators of hippocampal activity and indeed CNS activity increases, it is becoming clear that these hormones play a vital role.

It is clear from the above data that all three hormones discussed can modulate hippocampal function in similar ways. Given their closely associated functions it is likely that these three systems interact in the hippocampus although evidence of this is scant. In other systems a number of studies indicate that leptin and insulin signalling can converge at a number of points. Leptin reduces insulin-induced glucogenesis and tyrosine phosphorylation of IRS-1 by insulin.¹²⁵ Also, effects of insulin on hepatic glucose metabolism are enhanced by leptin.¹²⁶ A number of effects mediated by both these hormones involves modulation of PI3K signalling, which may act as a point of convergence for the different systems. In hypothalamic neurons for example, the activation of K_{ATP} channels by leptin is mimicked by insulin¹²⁷ and this activation is mediated by PI3-kinase in both cases.^{128,129}

Two major signalling systems, the PI3K/Akt and the Ras/Raf/MEK pathways can both be activated by either estrogens, insulin or leptin via activation of their respective receptor complexes (Fig. 1). PI3K and GSK3 β have both recently been shown to be involved in either AMPA receptor trafficking or synaptic plasticity in the

hippocampus^{67,130,131} so it is therefore possible that this signalling pathway could form a point of convergence for the modulatory effects ascribed to these three hormones. Furthermore, the Ras/Raf/MEK signalling cascade has also been shown to be involved in a number of forms of synaptic plasticity in the hippocampus¹³² and so could provide an additional point of integration for the actions of these three hormonal systems in hippocampal neurons.

What is clear is that a greater understanding of hormonal signalling and the interplay between different hormonal and neuronal systems is vital to understanding the complex mechanisms underlying processes such as synaptic plasticity and the modulation of dendritic morphology.

References

- Maffei M, Halaas J, Ravussin E, Pratley RE, Lee GH, Zhang Y, et al. Leptin levels in human and rodent: measurement of plasma leptin and ob RNA in obese and weight-reduced subjects. *Nat Med* 1995; 1:1155-61.
- Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med* 1996; 334:292-5.
- Morash B, Li A, Murphy PR, Wilkinson M, Ur E. Leptin gene expression in the brain and pituitary gland. *Endocrinology* 1999; 140:5995-8.
- Spiegelman BM, Flier JS. Obesity and the regulation of energy balance. *Cell* 2001; 104:531-43.
- Karsenty G. Convergence between bone and energy homeostases: leptin regulation of bone mass. *Cell Metab* 2006; 4:341-8.
- Cervero A, Dominguez F, Horcajadas JA, Quinonero A, Pellicer A, Simon C. The role of the leptin in reproduction. *Curr Opin Obstet Gynecol* 2006; 18:297-303.
- Malendowicz LK, Rucinski M, Belloni AS, Ziolkowska A, Nussdorfer GG. Leptin and the regulation of the hypothalamic-pituitary-adrenal axis. *Int Rev Cytol* 2007; 263:63-102.
- Harvey J, Shanley LJ, O'Malley D, Irving AJ. Leptin: a potential cognitive enhancer? *Biochem Soc Trans* 2005; 33:1029-32.
- Irving AJ, Wallace L, Durakoglugil D, Harvey J. Leptin enhances NR2B-mediated N-methyl-D-aspartate responses via a mitogen-activated protein kinase-dependent process in cerebellar granule cells. *Neuroscience* 2006; 138:1137-48.
- O'Malley D, MacDonald N, Mizelinska S, Connolly CN, Irving AJ, Harvey J. Leptin promotes rapid dynamic changes in hippocampal dendritic morphology. *Mol Cell Neurosci* 2007; 35:559-72.
- Tartaglia LA, Dembski M, Weng X, Deng N, Culpepper J, Devos R, et al. Identification and expression cloning of a leptin receptor, OB-R. *Cell* 1995; 83:1263-71.
- Tartaglia LA. The leptin receptor. *J Biol Chem* 1997; 272:6093-6.
- Uotani S, Bjorbaek C, Tornoe J, Flier JS. Functional properties of leptin receptor isoforms: internalization and degradation of leptin and ligand-induced receptor downregulation. *Diabetes* 1999; 48:279-86.

14. Hileman SM, Tornøe J, Flier JS, Bjorbaek C. Transcellular transport of leptin by the short leptin receptor isoform Ob-Ra in Madin-Darby Canine Kidney cells. *Endocrinology* 2000; 141:1955-61.
15. Banks WA, Niehoff ML, Martin D, Farrell CL. Leptin transport across the blood-brain barrier of the Koletsky rat is not mediated by a product of the leptin receptor gene. *Brain Res* 2002; 950:130-6.
16. DeVries SH, Schwartz EA. Kainate receptors mediate synaptic transmission between cones and 'Off' bipolar cells in a mammalian retina. *Nature* 1999; 397:157-60.
17. Savioz A, Charnay Y, Huguenin C, Graviou C, Greggio B, Bouras C. Expression of leptin receptor mRNA (long form splice variant) in the human cerebellum. *Neuroreport* 1997; 8:123-6.
18. Elmquist JK, Bjorbaek C, Ahima RS, Flier JS, Saper CB. Distributions of leptin receptor mRNA isoforms in the rat brain. *J Comp Neurol* 1998; 395:535-47.
19. Hakansson ML, Brown H, Ghilardi N, Skoda RC, Meister B. Leptin receptor immunoreactivity in chemically defined target neurons of the hypothalamus. *J Neurosci* 1998; 18:559-72.
20. Mercer JG, Hoggard N, Williams LM, Lawrence CB, Hannah LT, Trayhurn P. Localization of leptin receptor mRNA and the long form splice variant (Ob-Rb) in mouse hypothalamus and adjacent brain regions by in situ hybridization. *FEBS Lett* 1996; 387:113-6.
21. Figlewicz DP, Evans SB, Murphy J, Hoen M, Baskin DG. Expression of receptors for insulin and leptin in the ventral tegmental area/substantia nigra (VTA/SN) of the rat. *Brain Res* 2003; 964:107-15.
22. Huang XF, Koutcherov I, Lin S, Wang HQ, Storlien L. Localization of leptin receptor mRNA expression in mouse brain. *Neuroreport* 1996; 7:2635-8.
23. Shanley LJ, O'Malley D, Irving AJ, Ashford ML, Harvey J. Leptin inhibits epileptiform-like activity in rat hippocampal neurones via PI3-kinase-driven activation of BK channels. *J Physiol* 2002; 545:933-44.
24. Bliss TV, Collingridge GL. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 1993; 361:31-9.
25. Bear MF, Abraham WC. Long-term depression in hippocampus. *Annu Rev Neurosci* 1996; 19:437-62.
26. Li XL, Aou S, Oomura Y, Hori N, Fukunaga K, Hori T. Impairment of long-term potentiation and spatial memory in leptin receptor-deficient rodents. *Neuroscience* 2002; 113:607-15.
27. Winocur G, Greenwood CE, Piroli GG, Grillo CA, Reznikov LR, Reagan LP, et al. Memory impairment in obese Zucker rats: an investigation of cognitive function in an animal model of insulin resistance and obesity. *Behav Neurosci* 2005; 119:1389-95.
28. Farr SA, Banks WA, Morley JE. Effects of leptin on memory processing. *Peptides* 2006; 27:1420-5.
29. Oomura Y, Hori N, Shiraishi T, Fukunaga K, Takeda H, Tsuji M, et al. Leptin facilitates learning and memory performance and enhances hippocampal CA1 long-term potentiation and CaMK II phosphorylation in rats. *Peptides* 2006; 27:2738-49.
30. Wayner MJ, Armstrong DL, Phelix CF, Oomura Y. Orexin-A (Hypocretin-1) and leptin enhance LTP in the dentate gyrus of rats in vivo. *Peptides* 2004; 25:991-6.
31. Shanley LJ, Irving AJ, Harvey J. Leptin enhances NMDA receptor function and modulates hippocampal synaptic plasticity. *J Neurosci* 2001; 21:186.
32. Yuste R, Bonhoeffer T. Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annu Rev Neurosci* 2001; 24:1071-89.
33. Romano C, Sesma MA, McDonald CT, O'Malley K, Van den Pol AN, Olney JW. Distribution of metabotropic glutamate receptor mGluR5 immunoreactivity in rat brain. *J Comp Neurol* 1995; 355:455-69.
34. Barria A, Malinow R. Subunit-specific NMDA receptor trafficking to synapses. *Neuron* 2002; 35:345-53.
35. Tovar KR, Westbrook GL. The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses in vitro. *J Neurosci* 1999; 19:4180-8.
36. Ahmadian G, Ju W, Liu L, Wyszynski M, Lee SH, Dunah AW, et al. Tyrosine phosphorylation of GluR2 is required for insulin-stimulated AMPA receptor endocytosis and LTD. *Embo J* 2004; 23:1040-50.
37. Massey PV, Johnson BE, Moulton PR, Auberson YP, Brown MW, Molnar E, et al. Differential roles of NR2A and NR2B-containing NMDA receptors in cortical long-term potentiation and long-term depression. *J Neurosci* 2004; 24:7821-8.
38. Durakoglugil M, Irving AJ, Harvey J. Leptin induces a novel form of NMDA receptor-dependent long-term depression. *J Neurochem* 2005; 95:396-405.
39. Havrankova J, Roth J, Brownstein M. Insulin receptors are widely distributed in the central nervous system of the rat. *Nature* 1978; 272:827-9.
40. Havrankova J, Brownstein M, Roth J. Insulin and insulin receptors in rodent brain. *Diabetologia* 1981; 20:268-73.
41. Havrankova J, Roth J, Brownstein MJ. Insulin receptors in brain. *Adv Metab Disord* 1983; 10:259-68.
42. Wozniak M, Rydzewski B, Baker SP, Raizada MK. The cellular and physiological actions of insulin in the central nervous system. *Neurochem Int* 1993; 22:1-10.
43. Abbott MA, Wells DG, Fallon JR. The insulin receptor tyrosine kinase substrate p58/53 and the insulin receptor are components of CNS synapses. *J Neurosci* 1999; 19:7300-8.
44. Banks WA. The source of cerebral insulin. *Eur J Pharmacol* 2004; 490:5-12.
45. Clarke DW, Mudd L, Boyd FT Jr, Fields M, Raizada MK. Insulin is released from rat brain neuronal cells in culture. *J Neurochem* 1986; 47:831-6.
46. Schechter R, Whitmire J, Wheat GS, Beju D, Jackson KW, Harlow R, et al. 3rd. Immunohistochemical and in situ hybridization study of an insulin-like substance in fetal neuron cell cultures. *Brain Res* 1994; 636:9-27.
47. Devaskar SU, Giddings SJ, Rajakumar PA, Carnaghi LR, Menon RK, Zahm DS. Insulin gene expression and insulin synthesis in mammalian neuronal cells. *J Biol Chem* 1994; 269:8445-54.
48. Niswender KD, Schwartz MW. Insulin and leptin revisited: adiposity signals with overlapping physiological and intracellular signaling capabilities. *Front Neuroendocrinol* 2003; 24:1-10.
49. Wang YT, Linden DJ. Expression of cerebellar long-term depression requires postsynaptic clathrin-mediated endocytosis. *Neuron* 2000; 25:635-47.
50. Wan Q, Xiong ZG, Man HY, Ackerley CA, Brauton J, Lu WY, et al. Recruitment of functional GABA(A) receptors to postsynaptic domains by insulin. *Nature* 1997; 388:686-90.
51. Skeberdis VA, Lan J, Zheng X, Zukin RS, Bennett MV. Insulin promotes rapid delivery of N-methyl-D-aspartate receptors to the cell surface by exocytosis. *Proc Natl Acad Sci USA* 2001; 98:3561-6.
52. van der Heide LP, Ramakers GM, Smidt MP. Insulin signaling in the central nervous system: learning to survive. *Prog Neurobiol* 2006; 79:205-21.
53. Craft S, Peskind E, Schwartz MW, Schellenberg GD, Raskind M, Porte D Jr. Cerebrospinal fluid and plasma insulin levels in Alzheimer's disease: relationship to severity of dementia and apolipoprotein E genotype. *Neurology* 1998; 50:164-8.
54. White MF. The insulin signalling system and the IRS proteins. *Diabetologia* 1997; 40:2-17.
55. Taha C, Klip A. The insulin signaling pathway. *J Membr Biol* 1999; 169:1-12.
56. van der Heide LP, Hoekman MF, Biessels GJ, Gispen WH. Insulin inhibits extracellular regulated kinase 1/2 phosphorylation in a phosphatidylinositol 3-kinase (PI3) kinase-dependent manner in Neuro2a cells. *J Neurochem* 2003; 86:86-91.
57. White MF. The IRS-signalling system: a network of docking proteins that mediate insulin action. *Mol Cell Biochem* 1998; 182:3-11.
58. Wu SP, Lu KT, Chang WC, Gean PW. Involvement of mitogen-activated protein kinase in hippocampal long-term potentiation. *J Biomed Sci* 1999; 6:409-17.
59. Zhao W, Chen H, Xu H, Moore E, Meiri N, Quon MJ, et al. Brain insulin receptors and spatial memory. Correlated changes in gene expression, tyrosine phosphorylation, and signaling molecules in the hippocampus of water maze trained rats. *J Biol Chem* 1999; 274:34893-902.
60. Izquierdo LA, Barros DM, Ardenghi PG, Pereira P, Rodrigues C, Choi H, et al. Different hippocampal molecular requirements for short- and long-term retrieval of one-trial avoidance learning. *Behav Brain Res* 2000; 111:93-8.
61. Hodgekiss JP, Kelly JS. Only 'de novo' long-term depression (LTD) in the rat hippocampus in vitro is blocked by the same low concentration of FK506 that blocks LTD in the visual cortex. *Brain Res* 1995; 705:241-46.
62. Sanna PP, Cammalleri M, Berton F, Simpson C, Lütjens R, Bloom FE, Francesconi W. Phosphatidylinositol 3-kinase is required for the expression but not for the induction or the maintenance of long-term potentiation in the hippocampal CA1 region. *J Neurosci* 2002; 22:3359-65.
63. Hou L, Klann E. Activation of the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway is required for metabotropic glutamate receptor-dependent long-term depression. *J Neurosci* 2004; 24:6352-61.
64. Man HY, Lin JW, Ju WH, Ahmadian G, Liu L, Becker LE, et al. Regulation of AMPA receptor-mediated synaptic transmission by clathrin-dependent receptor internalization. *Neuron* 2000; 25:649-62.
65. Huang CC, Lee CC, Hsu KS. An investigation into signal transduction mechanisms involved in insulin-induced long-term depression in the CA1 region of the hippocampus. *J Neurochem* 2004; 89:217-31.
66. van der Heide LP, Kamal A, Artola A, Gispen WH, Ramakers GM. Insulin modulates hippocampal activity-dependent synaptic plasticity in a N-methyl-D-aspartate receptor and phosphatidylinositol-3-kinase-dependent manner. *J Neurochem* 2005; 94:1158-66.
67. Man HY, Wang Q, Lu WY, Ju WH, Ahmadian G, Liu L, et al. Activation of PI3-kinase is required for AMPA receptor insertion during LTP of mEPSCs in cultured hippocampal neurons. *Neuron* 2003; 38:611-24.
68. Passafaro M, Piech V, Sheng M. Subunit-specific temporal and spatial patterns of AMPA receptor exocytosis in hippocampal neurons. *Nat Neurosci* 2001; 4:917-26.
69. Collingridge GL, Isaac JT, Wang YT. Receptor trafficking and synaptic plasticity. *Nat Rev Neurosci* 2004; 5:952-62.
70. Liu L, Brown JC, 3rd, Webster WW, Morrisett RA, Monaghan DT. Insulin potentiates N-methyl-D-aspartate receptor activity in *Xenopus* oocytes and rat hippocampus. *Neurosci Lett* 1995; 192:5-8.
71. Liao GY, Leonard JP. Insulin modulation of cloned mouse NMDA receptor currents in *Xenopus* oocytes. *J Neurochem* 1999; 73:1510-9.
72. Lee CC, Huang CC, Wu MY, Hsu KS. Insulin stimulates postsynaptic density-95 protein translation via the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway. *J Biol Chem* 2005; 280:18543-50.
73. Sheng M. The postsynaptic NMDA-receptor-PSD-95 signaling complex in excitatory synapses of the brain. *J Cell Sci* 2001; 114:1251.
74. Niblock MM, Bruno-Bechtold JK, Riddle DR. Insulin-like growth factor I stimulates dendritic growth in primary somatosensory cortex. *J Neurosci* 2000; 20:4165-76.
75. Cheng CM, Mervis RF, Niu SL, Salem N Jr, Witters LA, Tseng V, et al. Insulin-like growth factor 1 is essential for normal dendritic growth. *J Neurosci Res* 2003; 73:1-9.
76. Schulingkamp RJ, Pagano TC, Hung D, Raffa RB. Insulin receptors and insulin action in the brain: review and clinical implications. *Neurosci Biobehav Rev* 2000; 24:855-72.

77. McEwen B. Estrogen actions throughout the brain. *Recent Prog Horm Res* 2002; 57:357-84.
78. Vasudevan N, Pfaff DW. Membrane-initiated actions of estrogens in neuroendocrinology: emerging principles. *Endocr Rev* 2007; 28:1-19.
79. Levin ER. Integration of the extranuclear and nuclear actions of estrogen. *Mol Endocrinol* 2005; 19:1951-9.
80. Weiland NG, Orikasa C, Hayashi S, McEwen BS. Distribution and hormone regulation of estrogen receptor immunoreactive cells in the hippocampus of male and female rats. *J Comp Neurol* 1997; 388:603-12.
81. Mitra SW, Hoskin E, Yudkovitz J, Pear L, Wilkinson HA, Hayashi S, et al. Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* 2003; 144:2055-67.
82. Register TC, Shively CA, Lewis CE. Expression of estrogen receptor alpha and beta transcripts in female monkey hippocampus and hypothalamus. *Brain Res* 1998; 788:320-2.
83. Simpkins JW, Wang J, Wang X, Perez E, Prokai L, Dykens JA. Mitochondria play a central role in estrogen-induced neuroprotection. *Curr Drug Targets CNS Neurol Disord* 2005; 4:69-83.
84. Toran-Allerand CD. Minireview: A plethora of estrogen receptors in the brain: where will it end? *Endocrinology* 2004; 145:1069-74.
85. Prange-Kiel J, Wehrenberg U, Jarry H, Rune GM. Para/autocrine regulation of estrogen receptors in hippocampal neurons. *Hippocampus* 2003; 13:226-34.
86. Rune GM, Frotscher M. Neurosteroid synthesis in the hippocampus: role in synaptic plasticity. *Neuroscience* 2005; 136:833-42.
87. Kawato S, Hojo Y, Kimoto T. Histological and metabolism analysis of P450 expression in the brain. *Methods Enzymol* 2002; 357:241-9.
88. Hojo Y, Hattori TA, Enami T, Furukawa A, Suzuki K, Ishii HT, et al. Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. *Proc Natl Acad Sci USA* 2004; 101:865-70.
89. Spencer JL, Waters EM, Romeo RD, Wood GE, Milner TA, McEwen BS. Uncovering the mechanisms of estrogen effects on hippocampal function. *Front Neuroendocrinol* 2007.
90. Milner TA, Ayoola K, Drake CT, Herrick SR, Tabori NE, McEwen BS, et al. Ultrastructural localization of estrogen receptor beta immunoreactivity in the rat hippocampal formation. *J Comp Neurol* 2005; 491:81-95.
91. Shughrue PJ, Merchenthaler I. Distribution of estrogen receptor beta immunoreactivity in the rat central nervous system. *J Comp Neurol* 2001; 436:64-81.
92. Gould E, Woolley CS, Frankfurt M, McEwen BS. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 1990; 10:1286-91.
93. Bleakman D, Ballyk BA, Schoepp DD, Palmer AJ, Bath CP, Sharpe EF, et al. Activity of 2,3-benzodiazepines at native rat and recombinant human glutamate receptors in vitro: stereospecificity and selectivity profiles. *Neuropharmacology* 1996; 35:1689-702.
94. Murphy DD, Segal M. Regulation of dendritic spine density in cultured rat hippocampal neurons by steroid hormones. *J Neurosci* 1996; 16:4059-68.
95. Gonzalez-Burgos I, Alejandro-Gomez M, Cervantes M. Spine-type densities of hippocampal CA1 neurons vary in proestrus and estrus rats. *Neurosci Lett* 2005; 379:52-4.
96. Tada T, Sheng M. Molecular mechanisms of dendritic spine morphogenesis. *Curr Opin Neurobiol* 2006; 16:95-101.
97. Rapp SR, Espeland MA, Shumaker SA, Henderson VW, Brunner RL, Manson JE, et al. Effect of estrogen plus progestin on global cognitive function in postmenopausal women: the Women's Health Initiative Memory Study: a randomized controlled trial. *JAMA* 2003; 289:2663-72.
98. Lee SJ, Romeo RD, Svenningsson P, Campomanes CR, Allen PB, Greengard P, et al. Estradiol affects spinophilin protein differently in gonadectomized males and females. *Neuroscience* 2004; 127:983-8.
99. Brakeman PR, Lanahan AA, O'Brien R, Roche K, Barnes CA, Hagan RL, et al. Homer: a protein that selectively binds metabotropic glutamate receptors. *Nature* 1997; 386:284-8.
100. Zhao L, Chen S, Ming Wang J, Brinton RD. 17beta-estradiol induces Ca²⁺ influx, dendritic and nuclear Ca²⁺ rise and subsequent cyclic AMP response element-binding protein activation in hippocampal neurons: a potential initiation mechanism for estrogen neurotrophism. *Neuroscience* 2005; 132:299-311.
101. Akama KT, McEwen BS. Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway. *J Neurosci* 2003; 23:2333-9.
102. Teyler TJ, Vardaris RM, Lewis D, Rawitch AB. Gonadal steroids: effects on excitability of hippocampal pyramidal cells. *Science* 1980; 209:1017-8.
103. Foy MR, Xu J, Xie X, Brinton RD, Thompson RF, Berger TW. 17beta-estradiol enhances NMDA receptor-mediated EPSPs and long-term potentiation. *J Neurophysiol* 1999; 81:925-9.
104. O'Neill MJ, Bogaert L, Hicks CA, Bond A, Ward MA, Ebinger G, et al. LY377770, a novel iGlu5 kainate receptor antagonist with neuroprotective effects in global and focal cerebral ischemia. *Neuropharmacology* 2000; 39:1575-88.
105. Fugger HN, Kumar A, Lubahn DB, Korach KS, Foster TC. Examination of estradiol effects on the rapid estradiol mediated increase in hippocampal synaptic transmission in estrogen receptor alpha knockout mice. *Neurosci Lett* 2001; 309:207-9.
106. Kim MT, Soussou W, Gholmieh G, Ahuja A, Tanguay A, Berger TW, et al. 17beta-Estradiol potentiates field excitatory postsynaptic potentials within each subfield of the hippocampus with greatest potentiation of the associational/commissural afferents of CA3. *Neuroscience* 2006; 141:391-406.
107. Wong M, Moss RL. Long-term and short-term electrophysiological effects of estrogen on the synaptic properties of hippocampal CA1 neurons. *J Neurosci* 1992; 12:3217-25.
108. Rudick CN, Woolley CS. Selective estrogen receptor modulators regulate phasic activation of hippocampal CA1 pyramidal cells by estrogen. *Endocrinology* 2003; 144:179-87.
109. Warren SG, Humphreys AG, Juraska JM, Greenough WT. LTP varies across the estrous cycle: enhanced synaptic plasticity in proestrus rats. *Brain Res* 1995; 703:26-30.
110. Cordoba Montoya DA, Carrer HF. Estrogen facilitates induction of long term potentiation in the hippocampus of awake rats. *Brain Res* 1997; 778:430-8.
111. Wu Z, Desmond NL, Levy WB. Homosynaptic long-term depression of CA3-CA3 synapses in the in vivo hippocampus. *Brain Res* 1998; 789:335-8.
112. Zamani MR, Desmond NL, Levy WB. Estradiol modulates long-term synaptic depression in female rat hippocampus. *J Neurophysiol* 2000; 84:1800-8.
113. Ito K, Skinkle KL, Hicks TP. Age-dependent, steroid-specific effects of oestrogen on long-term potentiation in rat hippocampal slices. *J Physiol* 1999; 515:209-20.
114. Whitlock JR, Heynen AJ, Shuler MG, Bear MF. Learning induces long-term potentiation in the hippocampus. *Science* 2006; 313:1093-7.
115. Norris CM, Korol DL, Foster TC. Increased susceptibility to induction of long-term depression and long-term potentiation reversal during aging. *J Neurosci* 1996; 16:5382-92.
116. Sandstrom NJ, Williams CL. Spatial memory retention is enhanced by acute and continuous estradiol replacement. *Horm Behav* 2004; 45:128-35.
117. Xu X, Zhang Z. Effects of estradiol benzoate on learning-memory behavior and synaptic structure in ovariectomized mice. *Life Sci* 2006; 79:1553-60.
118. Li C, Brake WG, Romeo RD, Dunlop JC, Gordon M, Buzescu R, et al. Estrogen alters hippocampal dendritic spine shape and enhances synaptic protein immunoreactivity and spatial memory in female mice. *Proc Natl Acad Sci USA* 2004; 101:2185-90.
119. Loy R, Gerlach JL, McEwen BS. Autoradiographic localization of estradiol-binding neurons in the rat hippocampal formation and entorhinal cortex. *Brain Res* 1988; 467:245-51.
120. Azcoitia I, Sierra A, Garcia-Segura LM. Localization of estrogen receptor beta-immunoreactivity in astrocytes of the adult rat brain. *Glia* 1999; 26:260-7.
121. Shughrue PJ, Scrimo PJ, Merchenthaler I. Estrogen binding and estrogen receptor characterization (ERalpha and ERbeta) in the cholinergic neurons of the rat basal forebrain. *Neuroscience* 2000; 96:41-9.
122. Mannel P, Brinton RD. Estrogen receptor protein interaction with phosphatidylinositol 3-kinase leads to activation of phosphorylated Akt and extracellular signal-regulated kinase 1/2 in the same population of cortical neurons: a unified mechanism of estrogen action. *J Neurosci* 2006; 26:9439-47.
123. Milner TA, McEwen BS, Hayashi S, Li CJ, Reagan LP, Alves SE. Ultrastructural evidence that hippocampal alpha estrogen receptors are located at extranuclear sites. *J Comp Neurol* 2001; 429:355-71.
124. Liu F, Day M, Muniz LC, Bitran D, Arias R, Revilla-Sanchez R, et al. Activation of estrogen receptor-beta regulates hippocampal synaptic plasticity and improves memory. *Nat Neurosci* 2008; 11:334-43.
125. Cohen B, Novick D, Rubinstein M. Modulation of insulin activities by leptin. *Science* 1996; 274:1185-8.
126. Aiston S, Agius L. Leptin enhances glycogen storage in hepatocytes by inhibition of phosphorylase and exerts an additive effect with insulin. *Diabetes* 1999; 48:15-20.
127. Spanwick D, Smith MA, Mirshamsi S, Routh VH, Ashford ML. Insulin activates ATP-sensitive K⁺ channels in hypothalamic neurons of lean, but not obese rats. *Nat Neurosci* 2000; 3:757-8.
128. Harvey J, Hardy SC, Irving AJ, Ashford ML. Leptin activation of ATP-sensitive K⁺ (KATP) channels in rat CRI-G1 insulinoma cells involves disruption of the actin cytoskeleton. *J Physiol* 2000; 527:95-107.
129. Mirshamsi S, Laidlaw HA, Ning K, Anderson E, Burgess LA, Gray A, et al. Leptin and insulin stimulation of signalling pathways in arcuate nucleus neurones: PI3K dependent actin reorganization and K_{ATP} channel activation. *BMC Neurosci* 2004; 5:54.
130. Baxter AW, Wyllie DJ. Phosphatidylinositol 3 kinase activation and AMPA receptor subunit trafficking underlie the potentiation of miniature EPSC amplitudes triggered by the activation of L-type calcium channels. *J Neurosci* 2006; 26:5456-69.
131. Peineau S, Taghibiglou C, Bradley C, Wong TP, Liu L, Lu J, et al. LTP inhibits LTD in the hippocampus via regulation of GSK3beta. *Neuron* 2007; 53:703-17.
132. Sweatt JD. The neuronal MAP kinase cascade: a biochemical signal integration system subserving synaptic plasticity and memory. *J Neurochem* 2001; 76:1-10.